

IN THE CLAIMS:

*Please cancel claims 1-8, 18, and 19. A complete listing of claims now pending appears below:*

Claims 1-8 (cancelled).

~~9.~~<sup>1</sup> (original) A method for preparing an organometallic complex useful for olefin polymerization, said method comprising:

- (a) converting a pentacyclic diketone to a triquinane diene;
  - (b) doubly deprotonating the triquinane diene to produce a triquinane dianion;
- and

(c) reacting the dianion with a transition metal source to give an organometallic complex that incorporates a chelating, dianionic triquinane ligand.

~~10.~~<sup>2</sup> (original) The method of claim ~~9~~<sup>1</sup> wherein the pentacyclic diketone is produced by (a) reacting a cyclopentadiene and a p-benzoquinone to produce a Diels-Alder adduct; and (b) photolyzing the Diels-Alder adduct to effect a [2+2] cycloaddition reaction to give the pentacyclic diketone.

~~11.~~<sup>3</sup> (original) The method of claim ~~9~~<sup>1</sup> wherein step (a) is accomplished by first heating the pentacyclic diketone to cause a [2+2] cycloreversion reaction to give a *cis,syn,cis*-triquinane bis(enone), followed by conversion of the bis(enone) to the triquinane diene.

~~12.~~<sup>4</sup> (original) The method of claim ~~11~~<sup>3</sup> wherein the bis(enone) is converted to the triquinane diene by (a) reacting the bis(enone) with an arylhydrazine to produce an arylhydrazone; and (b) reducing the arylhydrazone to the diene by reacting it with an alkali metal cyanoborohydride or catecholborane.

~~13.~~<sup>5</sup> (original) The method of claim ~~11~~<sup>3</sup> wherein the bis(enone) is converted to the triquinane diene by reacting it with a trialkylhydrosilane in the presence of a Lewis acid.

~~14.~~<sup>6</sup> (original) The method of claim ~~9~~<sup>1</sup> wherein step (a) is accomplished by first converting the pentacyclic diketone to a pentacyclic hydrocarbon by reducing the carbonyl groups to methylene groups, and then heating the pentacyclic hydrocarbon to cause a [2+2] cycloreversion reaction to give the triquinane diene.

~~15.~~<sup>7</sup> (original) The method of claim ~~9~~<sup>1</sup> wherein the pentacyclic diketone is homologated by reacting it with diazomethane prior to conversion to the triquinane diene.

~~16.~~<sup>8</sup> (original) A method for preparing an organometallic complex useful for olefin polymerization, said method comprising:

(a) reacting a cyclopentadiene and a p-benzoquinone to produce a Diels-Alder adduct;

(b) photolyzing the Diels-Alder adduct to effect a [2+2] cycloaddition reaction to give a pentacyclic diketone;

(c) converting the pentacyclic diketone to a triquinane diene;

(d) doubly deprotonating the triquinane diene to produce a triquinane dianion;  
and

(e) reacting the dianion with a transition metal source to give an organometallic complex that incorporates a chelating, dianionic triquinane ligand.

~~17.~~<sup>9</sup> (original) The method of claim ~~16~~<sup>8</sup> wherein the Diels-Alder adduct is produced from cyclopentadiene and p-benzoquinone.

Claims 18-19 (cancelled).